

[psng]mm  
 [1] 1 [1] 1 [1] 1 textbf{cmbxti10} textbf{fss} cmssbx10 mathbf{cmbxti10} mathbf{fss} cmssbx10 upmath  
 eurm10 AMSa msam10 0upmath19 0upmath16 0upmath40 3AMSa36 3AMSa3E  
 normalcmrmit boldcmrbxit [1] 1 [1] 1 [1] 1 normalcmrbxit boldcmrbxit normalcmssbxn bold-  
 cmssbxn @mathgroup@group @mathgroup@normal@groupeurmn @mathgroup@bold@groupeurn @math-  
 group@group @mathgroup@normal@groupmsamn @mathgroup@bold@groupmsamn @group19 @group16  
 @group40 @group36 @group3E  
 [1] 1 [1] 1 [1] 1 OT1cmrbxit boldOT1cmrbxit OT1cmssbxn boldOT1cmssbxn UPMUeurmn UPM-  
 boldUeurbn AMSaUmsamn  $\Gamma 0UPM''19\Gamma 0UPM''16\Gamma 0UPM''40\Lambda 3AMSa''36\Lambda 3AMSa''3E$

The Pattern Speed of the OH/IR Stars in the Milky Way [Victor P. Debattista, Ortwin Gerhard and Maartje N. Sevenster] Victor P. Debattista<sup>1</sup>, Ortwin Gerhard<sup>1</sup> and Maartje N. Sevenster<sup>2</sup>

firstpage–lastpage 2001

document

firstpage

abstract We show how the continuity equation can be used to determine pattern speeds in the Milky Way Galaxy (MWG). This method, first discussed by Tremaine & Weinberg in the context of external galaxies, requires projected positions,  $(l, b)$ , and line-of-sight velocities for a spatially complete sample of relaxed tracers. If the local standard of rest (LSR) has a zero velocity in the radial direction ( $u_{\text{LSR}}$ ), then the quantity that is measured is  $\Delta V \equiv \Omega_p R_0 - V_{\text{LSR}}$ , where  $\Omega_p$  is the pattern speed of the non-axisymmetric feature,  $R_0$  is the distance of the Sun from the Galactic centre and  $V_{\text{LSR}}$  is the tangential motion of the LSR, including the circular velocity. We use simple models to assess the reliability of the method for measuring a single, constant pattern speed of either a bar or spiral in the inner MWG. We then apply the method to the OH/IR stars in the ATCA/VLA OH 1612 MHz survey of Sevenster *et al.*, finding  $\Delta V = 252 \pm 41 \text{ km s}^{-1}$ , if  $u_{\text{LSR}} = 0$ . Assuming further that  $R_0 = 8 \text{ kpc}$  and  $V_{\text{LSR}} = 220 \text{ km s}^{-1}$ , this gives  $\Omega_p = 59 \pm 5 \text{ km s}^{-1} \text{ kpc}^{-1}$  with a possible systematic error of perhaps  $10 \text{ km s}^{-1} \text{ kpc}^{-1}$ . The non-axisymmetric feature for which we measure this pattern speed must be in the disc of the MWG.